

On a new type of Double Sequences of Fuzzy Numbers.

Hemen Dutta¹ and B. Surender Reddy²

¹Department of Mathematics, Gauhati University, Kokrajhar Campus, Assam, India.

²Department of Mathematics, PGCS, Saifabad, Osmania University, Hyderabad, 500004, AP, India.

E-mail: hemen_dutta08@rediffmail.com
bsrmathou@yahoo.com

ABSTRACT

In this article we introduce the notion of Δ -statistically pre-Cauchy double sequence of fuzzy numbers and establish a criterion for arbitrary double sequence of fuzzy numbers to be Δ -statistically pre-Cauchy.

(Keywords: double sequence of fuzzy numbers, difference sequence, statistically pre-Cauchy double sequence, Orlicz function)

(AMS Subject Classification No: 46E30, 40A05, 40D25, 26A03, 46A45, 46A20)

INTRODUCTION

Fuzzy set theory, compared to other mathematical theories, is perhaps the most easily adaptable theory to practice. The main reason is that a fuzzy set has the property of relativity, variability, and inexactness in the definition of its elements. Instead of defining an entity in calculus by assuming that its role is exactly known, we can use fuzzy sets to define the same entity by allowing possible deviations and inexactness in its role. This representation suits well the uncertainties encountered in practical life, which make fuzzy sets a valuable mathematical tool. The concepts of fuzzy sets and fuzzy set operations were first introduced by Zadeh [35] and subsequently several authors have discussed various aspects of the theory and applications of fuzzy sets such as fuzzy topological spaces, similarity relations and fuzzy orderings, fuzzy measures of fuzzy events, fuzzy mathematical programming. Matloka [20] introduced bounded and convergent sequences of fuzzy numbers and studied their some properties. Later on sequences of fuzzy numbers have been discussed by Diamond and Kloeden [15], Nanda [21], Esi [13], Dutta [7, 8, 9], and many others.

A fuzzy number is a fuzzy set on the real axis, i.e., a mapping $u: R \rightarrow [0,1]$ which satisfies the following four conditions:

(i) u is normal, i.e., there exist an $x_0 \in R$ such that $u(x_0) = 1$.

(ii) u is fuzzy convex, i.e.

$u[\lambda x + (1-\lambda)y] \geq \min\{u(x), u(y)\}$ for all $x, y \in R$ and for all $\lambda \in [0,1]$.

(iii) u is upper semi-continuous.

(iv) The set $u_0 = \overline{\{x \in R : u(x) > 0\}}$ is compact,

where $\overline{\{x \in R : u(x) > 0\}}$ denotes the closure of

the set $\{x \in R : u(x) > 0\}$ in the usual topology of R .

We denote the set of all fuzzy numbers on R by E^1 and called it as the space of fuzzy numbers.

λ -level set u_λ of $u \in E^1$ is defined by:

$$[u]_\lambda = \{t \in R : u(t) \geq \lambda\} \text{ if } (0 < \lambda \leq 1), \\ = \overline{\{t \in R : u(t) > \lambda\}} \text{ if } (\lambda = 0)$$

The set u_λ is a closed, bounded and non-empty interval for each $\lambda \in (0,1)$ which is defined

by $u_\lambda = [u^-(\lambda), u^+(\lambda)]$. R can be embedded in E^1 , since each $r \in R$ can be regarded as a fuzzy number,

$$\overline{r(t)} = 1 \text{ if } t = r \\ = 0 \text{ if } t \neq r$$

Let W be the set of all closed bounded intervals A of real numbers such that $A = [A_1, A_2]$. Define the relation d on W as follows:

$$d(A, B) = \max\{|A_1 - B_1|, |A_2 - B_2|\}.$$

Then W, d is a complete metric space (See Diamond and Kloeden [14], Nanda [21]).

A fuzzy double sequence is a double infinite array of fuzzy real numbers. We denote a fuzzy double sequence by a_{mn} , where a_{mn} are fuzzy real numbers for each $m, n \in N$. The initial works on double sequences of real or complex terms is found in Bromwich [1]. Hardy [18] introduced the notion of regular convergence for double sequences of real or complex terms. The works on double sequence was further investigated by Basarir and Solanacan [2], Moricz [23], Tripathy and Dutta [31], Tripathy and Sarma [32] and many others.

The concept of statistical convergence was first introduced by Fast [16] and also independently by Buck [3] and Schoenberg [28] for real and complex sequences. Further this concept was studied by Salat [27], Fridy [17], Cannor [4, 5] and many others.

A double sequence of fuzzy number (x_{kl}) is called Δ -statistically convergent to L if

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \left| \{(j, k) : d(\Delta x_{jk}, L) \geq \varepsilon, j \leq m, k \leq n\} \right| = 0.$$

where the vertical bars indicate the number of elements in the set.

Definition 1: A double sequence of fuzzy number (x_{kl}) is called Δ -statistically pre-Cauchy if for every $\varepsilon > 0$ there exist $p = p(\varepsilon)$ and $q(\varepsilon)$ such that

$$\lim_{m,n \rightarrow \infty} \frac{1}{m^2 n^2} \left| \{(j, k) : d(\Delta x_{jk}, \Delta x_{pq}) \geq \varepsilon, j \leq m, k \leq n\} \right| = 0$$

In fact, the first order difference operator Δ can be viewed as an infinite triangular matrix as follows:

$$\Delta = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & \dots \\ 0 & 1 & -1 & 0 & 0 & \dots \\ 0 & 0 & 1 & -1 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

The fuzzy double sequence by a_{mn} can be expressed as an infinite matrix of fuzzy numbers as follows:

$$a_{mn} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} & \dots \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} & \dots \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

Now for any fuzzy double sequence by $a = a_{mn}$, we have

$$\Delta a = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & \dots \\ 0 & 1 & -1 & 0 & 0 & \dots \\ 0 & 0 & 1 & -1 & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} & \dots \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} & \dots \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

$$= \begin{pmatrix} a_{11} - a_{21} & a_{12} - a_{22} & \dots & a_{1n} - a_{2n} & \dots \\ a_{21} - a_{31} & a_{22} - a_{32} & \dots & a_{2n} - a_{3n} & \dots \\ a_{31} - a_{41} & a_{32} - a_{42} & \dots & a_{3n} - a_{4n} & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}.$$

This approach of construction of difference sequences is useful to study some properties of the spaces of such sequences.

An Orlicz Function is a function $M : [0, \infty) \rightarrow [0, \infty)$ which is continuous, non decreasing and convex with $M(0) = 0$, $M(x) > 0$ for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$.

If convexity of M is replaced by $M(x+y) \leq M(x) + M(y)$, then it is called a Modulus function (see Maddox[22]). An Orlicz function may be bounded or unbounded.

For example, $M(x) = x^p$ ($0 < p \leq 1$) is unbounded and $M(x) = \frac{x}{x+1}$ is bounded.

Lindesstrauss and Tzafriri [19] used the idea of Orlicz sequence space and introduced the sequence space ℓ_M as follows:

$$l_M = \{x = (x_k) \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty,$$

for some $\rho > 0$

They proved that ℓ_M is a Banach space normed by:

$$\|x\|_M = \|(x_k)\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}$$

The space l_M is closely related to the space l_p , which is an Orlicz sequence space with $M(x) = x^p$ for $1 \leq p < \infty$. An Orlicz function M satisfies the Δ_2 -condition ($M \in \Delta_2$ for short) if there exist constant $k \geq 2$ and $u_0 > 0$ such that $M(2u) \leq kM(u)$ whenever $|u| \leq u_0$. Note that an Orlicz function satisfies the inequality.

$$M(\lambda x) \leq \lambda M(x) \text{ for all } \lambda \text{ with } 0 < \lambda < 1.$$

The study of Orlicz sequence spaces have been made recently by various authors (cf [10], [11], [12], [25], [33], [34]).

In [6] Connor, Fridy and Kline proved that a bounded sequence $x = (x_i)$ is statistically pre-Cauchy if and only if $\lim_n \sum_{j,i \leq n} |x_i - x_j| = 0$.

We establish the following criterion for arbitrary double sequence of fuzzy numbers to be statistically pre-Cauchy.

MAIN RESULTS

Theorem 1: Let $x = (x_{jk})$ be the double sequence of fuzzy number and let M be a bounded Orlicz function. Then x is Δ -statistically pre-Cauchy if and only if

$$\lim_{m,n} \frac{1}{m^2 n^2} \sum_{j,p \leq m} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) = 0,$$

for some $\rho > 0$.

Proof : Suppose

$$\lim_{m,n} \frac{1}{m^2 n^2} \sum_{j,p \leq m} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) = 0,$$

for some $\rho > 0$. For each $\varepsilon > 0$, $\rho > 0$ and $m, n \in N$, we have

$$\begin{aligned} & \frac{1}{m^2 n^2} \sum_{j,p \leq m} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &= \frac{1}{m^2 n^2} \sum_{j,p \leq m, d(\Delta x_{jk}, \Delta x_{pq}) < \varepsilon} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &+ \frac{1}{m^2 n^2} \sum_{j,p \leq m, d(\Delta x_{jk}, \Delta x_{pq}) \geq \varepsilon} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &\geq \frac{1}{m^2 n^2} \sum_{j,p \leq m, d(\Delta x_{jk}, \Delta x_{pq}) \geq \varepsilon} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &\geq M(\varepsilon) \left(\frac{1}{m^2 n^2} |\{(j,k) : d(\Delta x_{jk}, \Delta x_{pq}) \geq \varepsilon, j \leq m, k \leq n\}| \right) \\ &\geq 0. \end{aligned}$$

Now suppose that x is Δ -statistically pre-Cauchy and that ε has been given. Let $\varepsilon > 0$ be such that $M(\delta) < \frac{\varepsilon}{2}$. Since M is bounded, there exist

an integer B such that $M(x) < \frac{B}{2}$

for all $x \geq 0$. Note that, for each $n \in N$.

$$\begin{aligned} & \frac{1}{m^2 n^2} \sum_{j,p \leq m} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &= \frac{1}{m^2 n^2} \sum_{j,p \leq m, d(\Delta x_{jk}, \Delta x_{pq}) < \delta} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &+ \frac{1}{m^2 n^2} \sum_{j,p \leq m, d(\Delta x_{jk}, \Delta x_{pq}) \geq \delta} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \\ &\leq M(\delta) + \frac{1}{m^2 n^2} \sum_{j,p \leq m, d(\Delta x_{jk}, \Delta x_{pq}) \geq \delta} \sum_{k,q \leq n} M\left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho}\right) \end{aligned}$$

$$\leq \frac{\varepsilon}{2} + \frac{B}{2} \left(\frac{1}{m^2 n^2} |\{(j, k) : d(\Delta x_{jk}, \Delta x_{pq}) \geq \delta, j \leq m, k \leq n\}| \right)$$

$$\leq \varepsilon + B \left(\frac{1}{m^2 n^2} |\{(j, k) : d(\Delta x_{jk}, \Delta x_{pq}) \geq \delta, j \leq m, k \leq n\}| \right)$$

(1)

Since x is Δ -statistically pre-Cauchy, there is N such that R.H.S of equation (1) is less than ε for all $n \in N$. Hence

$$\lim_{m,n} \frac{1}{m^2 n^2} \sum_{j,p \leq m} \sum_{k,q \leq n} M \left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho} \right) = 0.$$

Theorem 2: Let $x = (x_{jk})$ be the double sequence of fuzzy number and let M be a bounded Orlicz function. Then x is Δ -statistically convergent to L if and only if

$$\lim_{m,n} \frac{1}{mn} \sum_{j=1}^m \sum_{k=1}^n M \left(\frac{d(\Delta x_{jk}, L)}{\rho} \right) = 0$$

Proof: Suppose that

$$\lim_{m,n} \frac{1}{mn} \sum_{j=1}^m \sum_{k=1}^n M \left(\frac{d(\Delta x_{jk}, L)}{\rho} \right) = 0$$

with an Orlicz function M , then x is Δ -statistically convergent to L (see[22]). Conversely suppose that x is Δ -statistically convergent to L . We can prove in the similar manner to Theorem 1 that

$$\lim_{m,n} \frac{1}{mn} \sum_{j=1}^m \sum_{k=1}^n M \left(\frac{d(\Delta x_{jk}, L)}{\rho} \right) = 0$$

using that M is an Orlicz function.

Corollary 1: Let $x = (x_{jk})$ be the double sequence of fuzzy number. Then x is Δ -statistically pre-Cauchy if and only if

$$\lim_{m,n} \frac{1}{m^2 n^2} \sum_{j=1}^m \sum_{k=1}^n d(\Delta x_{jk}, \Delta x_{pq}) = 0$$

Proof: Let $A = \sup_{j,k} d \Delta x_{jk}, \bar{0}$ and define

$$M(x) = \frac{(1+2A)x}{1+x}$$

Then

$$M \left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho} \right) \leq (1+2A)d(\Delta x_{jk}, \Delta x_{pq})$$

and

$$M \left(\frac{d(\Delta x_{jk}, \Delta x_{pq})}{\rho} \right) = (1+2A) \frac{d(\Delta x_{jk}, \Delta x_{pq})}{1+d(\Delta x_{jk}, \Delta x_{pq})}$$

$$\geq \frac{(1+2A)d(\Delta x_{jk}, \Delta x_{pq})}{1+d(\Delta x_{jk}, \Delta x_{pq})}$$

$$\geq \frac{(1+2A)d(\Delta x_{jk}, \Delta x_{pq})}{1+2A}$$

$$= d(\Delta x_{jk}, \Delta x_{pq})$$

Hence

$$\lim_{m,n} \frac{1}{m^2 n^2} \sum_{j=1}^m \sum_{k=1}^n d(x_{jk}, x_{pq}) = 0$$

if and only if

$$\lim_{m,n} \frac{1}{m^2 n^2} \sum_{j=1}^m \sum_{k=1}^n M \left(\frac{d(x_{jk}, x_{pq})}{\rho} \right) = 0$$

and an immediate application of Theorem 1 completes the proof.

Corollary 2: Let $x = (x_{jk})$ be the double sequence of fuzzy number. Then x is Δ -statistically convergent to L if and only if

$$\lim_{m,n} \frac{1}{mn} \sum_{j=1}^m \sum_{k=1}^n d(\Delta x_{jk}, L) = 0$$

Proof: Let $A = \sup_{j,k} d \Delta x_{jk}, \bar{0}$ and define

$$M(x) = \frac{(1+A+L)x}{1+x}.$$

We can prove in the similar manner as in the proof of Corollary 1.

REFERENCES

- Bromwich, T.J.I. 1965. *An Introduction to the Theory of Infinite Series*. Macmillan & Co.: New York.
- Basarir, M. and O. Solancan. 1999. "On Some Double Sequence Spaces". *J. Indian Acad. Math.* 21(2):193-200.
- Buck, R.C. 1953, "Generalized Asymptote Density". *Amer J. of Math.* 75: 335-346.
- Connor, J.S. 1988. "The Statistical and Strong P-Cesaro Convergence of Sequences". *Analysis.* 8: 47-63.
- Connor, J.S. 1989. "On Strong Matrix Summability with Respect to a Modulus and Statistical Convergence". *Canad. Math. Bull.* 32:194-198.
- Connor, J.S., J. Fridy, and J. Kline. 1994. "Statistically Pre-Cauchy Sequences". *Analysis.* 14:311-317.
- Dutta, H. 2010. "On Some Complete Metric Spaces of Strongly Summable Sequences of Fuzzy Numbers". *Rend. Semin. Mat. Univ. Politec. Torino*, 68. (In Press).
- Dutta, H. 2010, "On Some New Type of Summable and Statistically Convergence Difference Sequences of Fuzzy Numbers". *J. Fuzzy Math.* 18(4). (In Press)
- Dutta, H. 2009, "On some Isometric Spaces of C_0^F , C^F and ℓ_∞^F ". *Acta Univ. Apulensis*, 19:107-112.
- Dutta, H. 2009. "On Köthe-Toeplitz and Null Duals of Some Difference Sequence Spaces Defined by Orlicz Functions". *Eur. J. Pure Appl. Math.* 2(4):554-563.
- Esi, A. and M. Et. 2000. "Some New Sequence Spaces Defined by a Sequence of Orlicz Functions". *Indian J. Pure Appl. Math.* 31(8):967-972.
- Et, M. 2001. "On Some New Orlicz Sequences Spaces". *J. Analysis.* 9:21-28.
- Esi, A. 2006. "On Some New Paranormed Sequence Spaces of Fuzzy Numbers Defined by Orlicz Functions and Statistical Convergence". *Mathematical Modelling and Analysis.* 1(4):379-388.
- Diamond, P. and P. Kloeden. 1990. "Metric Spaces of Fuzzy Sets, Fuzzy Sets and Systems". 35:241-249.
- Diamond, P. and P. Kloeden. 1994, "Metric Spaces of Fuzzy Sets". *Theory and Applications*, World Scientific: Singapore.
- Fast, H. 1951. "Sur la convergence statistique". *Colloq. Math.* 2:241-244.
- Fridy, J.A. 1985. "On Statistical Convergence". *Analysis.* 5:301-313.
- Hardy, G.H. 1917. "On the Convergence of Certain Multiple Series". *Proc. Camb. Phil. Soc.* 19: 86-95.
- Lindenstrauss, J. and L. Tzafriri. 1971. "On Orlicz Sequence Spaces". *Israel J. Math.* 10:379-390.
- Matloka, M. 1986. "Sequences of Fuzzy Numbers". *BUSEFAL.* 28:28-37.
- Nanda, S. 1989. "On Sequences of Fuzzy Numbers". *Fuzzy Sets and Systems.* 33:123-126.
- Maddox, I.J. 1986. "Sequence Spaces Defined by Modulus". *Math. Proc. Camb. Soc.* 100:161-166.
- Moricz, F. 1991. "Extension of Spaces c and c_0 from Single to Double Sequences". *Acta Math. Hung.* 57(1-2):129-136.
- Moricz, F. and B.E. Rhoades. 1988. "Almost Convergence of Double Sequences and Strong Regularity of Summability Matrices". *Math. Proc. Camb. Phil. Soc.* 104:283-294.
- Parashar, S.D. and B. Choudhary. 1994. "Sequence Spaces Defined by Orlicz Functions". *Indian J. Pure Appl. Math.* 25(4):419-428.
- Talo, Ö. and F. Başar. 2009. "Determination of the Duals of Classical Sets of Sequences of Fuzzy Numbers and Related Matrix Transformations". *Computers and Mathematics with Applications.* 58:717-733.
- Salat, T. 1980. "On Statistically Convergent Sequences of Real Numbers". *Math. Soovaca.* 30:139-150.
- Schoenberg, I.J. 1959. "The Integrability of Certain Functions and Related Sumability Methods". *Amer. Math. Monthly.* 66:361-375.
- Sarma, B. 2005. "Studies on Some Vector Valued Sequence Spaces and Köthe-Toeplitz Duals". (Ph.D. Thesis).
- Tripathy, B.C. 2003. "Statistically Convergent Double Sequences". *Tamkang J. Math.* 34(3):231-237.

31. Tripathy, B.C. and A.J. Dutta, 2007, "On Fuzzy Real-Valued Double Sequence Space ${}^2\ell_F^p$ ". *Math. Comput. Modelling*. 46(9-10):1294-1299.
32. Tripathy, B.C. and B. Sarma. 2008. "Statistically Convergent Difference Double Sequence Spaces". *Acta Math. Sinica*. 24(5):737-742.
33. V.A. Khan and Q.M. Danish Lohani. 2007. "Statistically pre-Cauchy Sequences and Orlicz Functions". *Southeast Asian Bull. of Math*. 31:1107-1112.
34. Yan, Y. 2004. "An Interpolation Inequality in Orlicz Spaces". *Southeast Asian Bull. of Math*. 28:931-936.
35. Zadeh, L.A. 1965. "Fuzzy Sets". *Information and Control*. 8:338-353.

ABOUT THE AUTHORS

Prof. Hemen Dutta is an Assistant Professor of Mathematics, Gauhati University, Kokrajhar Campus, Assam, India. He has published/accepted more than 40 research papers in different peer-reviewed journals. He is a reviewer of different journals and serves as an Associate Editorial Board Member of the International Journal of Open Problems in Computer Science and Mathematics. His research interests are in the areas of Mathematical Analysis and Fuzzy Mathematics.

Prof. B. Surender Reddy is an Associate Professor of Mathematics, Department of Mathematics, Osmania University, Hyderabad, Andhra Pradesh, India. He has published/accepted more than 25 papers in different peer-reviewed journals. His research interests are in the areas of Functional Analysis, Operator Theory and its applications, Mathematical Analysis and Fuzzy Mathematics.

SUGGESTED CITATION

Dutta, H. and B.S. Reddy. 2010. "On a new type of Double Sequences of Fuzzy Numbers". *Pacific Journal of Science and Technology*. 11(2):254-259.



[Pacific Journal of Science and Technology](http://www.akamaiuniversity.us/PJST.htm)